arXiv:1607.02223v1 [math.AT] 8 Jul 2016

Periodic points on T - fiber bundles over the circle

Weslem L. Silva and Rafael M. Souza

July 11, 2016

1 Introduction

Let $f : M \to M$ be a map and $x \in M$, where M a compact manifold. The point x is called a periodic point of f if there exists $n \in \mathbb{N}$ such that $f^n(x) = x$, in this case x a periodic point of f of period n. The set of all $\{x \in M|$ x is periodic$\}$ is called the set of periodic points of f and is denoted by $P(f)$.

If M is a compact manifold then the Nielsen theory can be generalized to periodic points. Boju Jiang introduced (Chapter 3 in [1]) a Nielsen-type homotopy invariant $NF_n(f)$ being a lower bound of the number of n-periodic points, for each g homotopic to f; $Fix(g^n) \geq NF_n(f)$. In case $dim(M) \geq 3$, M compact PL- manifold, then any map $f : M \to M$ is homotopic to a map g satisfying $Fix(g^n) = NF_n(f)$, this was proved in [2].

Consider a fiber bundle $F \to M \xrightarrow{p} B$ where F, M, B are closed manifolds and $f : M \to M$ a fiber-preserving map over B. In natural way is to study periodic points of f on M, that is, given $n \in \mathbb{N}$ we want to study the set $\{x \in M|f^n(x) = x\}$. The our main question is; when f can be deformed by a fiberwise homotopy to a map $g : M \to M$ such that $Fix(g^n) = \emptyset$?

In this paper we study periodic points of f when the fiber is the torus T and the base is the circle S^1.

2 General problem

Let $F \to M \xrightarrow{p} B$ be a fibration and $f : M \to M$ a fiber-preserving map over B, where F, M, B are closed manifolds. Given $n \in \mathbb{N}$, from relation $p \circ f = p$, we obtain $p \circ f^n = p$, thus $f^n : M \to M$ is also a fiber-preserving map for each $n \in \mathbb{N}$. We want to know when f can be deformed by a fiberwise homotopy

to a map $g : M \to M$ such that $Fix(g^n) = \emptyset$. The the following lemma give us a necessary condition to a positive answer the question above.

Lemma 2.1. *Let $f : M \to M$ be a fiber-preserving map. If some k, where k divides n, the map $f^k : M \to M$ can not be deformed to a fixed point free map, by a fiberwise homotopy, then can not exists, $g \sim_B f$, such that $g^n : M \to M$ is a fixed point free map.*

Proof. In fact, suppose that exists $g \sim_B f$ such that $Fix(g^n) = \emptyset$. Since $Fix(g^k) \subset Fix(g^n)$ and $Fix(g^k) \neq \emptyset$ then we have a contradiction. □

Therefore, a necessary condition is that for all k, where k divides n, the map $f^k : M \to M$ must be deformed by a fiberwise homotopy to a fixed point free map over B.

Note that for each n the square of the following diagram is commutative;

$$\begin{array}{ccccccccc}
\ldots & \longrightarrow & \pi_1(F, x_0) & \xrightarrow{i_\#} & \pi_1(M, x_0) & \xrightarrow{p_\#} & \pi_1(B, p(x_0)) & \longrightarrow & 0 \\
 & & \downarrow{\scriptstyle (f^n|_F)_\#} & & \downarrow{\scriptstyle f^n_\#} & & \downarrow{\scriptstyle Id} & & \\
\ldots & \longrightarrow & \pi_1(F, f^n(x_0)) & \xrightarrow{i_\#} & \pi_1(M, f^n(x_0)) & \xrightarrow{p_\#} & \pi_1(B, p(x_0)) & \longrightarrow & 0
\end{array}$$

Since in our case all spaces are path-connected then we will represent the generators of the groups $\pi_1(M, f^n(x_0))$ for each n, with the same letters. The same thing we will do with $\pi_1(T, f^n(0))$.

Let $M \times_B M$ be the pullback of $p : M \to B$ by $p : M \to B$ and $p_i : M \times_B M \to M, i = 1, 2,$ the projections to the first and the second coordinates, respectively.

The inclusion $M \times_B M - \Delta \hookrightarrow M \times_B M$, where Δ is the diagonal in $M \times_B M$, is replaced by the fiber bundle $q : E_B(M) \to M \times_B M$, whose fiber is denoted by $\mathcal{F}$. We have $\pi_m(E_B(M)) \approx \pi_m(M \times_B M - \Delta)$ where $E_B(M) = \{(x, \omega) \in B \times A^I | i(x) = \omega(0)\}$, with $A = M \times_B M$, $B = M \times_B M - \Delta$ and q is given by $q(x, \omega) = \omega(1)$.

E. Fadell and S. Husseini in [4] studied the problem to deform the map f^n, for each $n \in \mathbb{N}$, to a fixed point free map. They supposed that $dim(F) \geq 3$ and that F, M, B are closed manifolds. The necessary and sufficient condition to deform f^n is given by the following theorem that the proof can be find in [4].

Theorem 2.1. *Given $n \in \mathbb{N}$, the map $f^n : M \to M$ is deformable to a fixed point free map if and only if there exists a lift $\sigma(n)$ in the following diagram;*

$$\begin{array}{ccccc} \mathcal{F} & & \mathcal{F} & & \\ \downarrow & & \downarrow & & \\ E_B(f^n) & \xrightarrow{\bar{q}_{f^n}} & E_B(M) & & \\ \downarrow q_{f^n} & \nearrow_{\sigma(n)} & \downarrow q & & \\ M & \xrightarrow{1} M & \xrightarrow{(1,f^n)} & M \times_B M & \end{array} \tag{1}$$

where $E_B(f^n) \to M$ is the fiber bundle induced from q by $(1, f^n)$.

In the Theorem 2.1 we have $\pi_{j-1}(\mathcal{F}) \cong \pi_j(M \times_B M, M \times_B M - \Delta) \cong \pi_j(F, F - x)$ where x is a point in F. In this situation, that is, $dim(F) \geq 3$ the classical obstruction was used to find a cross section.

When F is a surface with Euler characteristic ≤ 0 then by Proposition 1.6 from [6] we have necessary e sufficient conditions to deform f^n to a fixed point free map over B. The next proposition gives a relation between a geometric diagram and our problem.

Proposition 2.1. *Let $f : M \to M$ be a fiber-preserving map over B. Then there is a map g, $g \sim_B f$, such that $Fix(g^n) = \emptyset$ if and only if there is a map $h_n : M \to M \times_B M - \Delta$ of the form $h_n = (Id, s^n)$, where $s : M \to M$, is fiberwise homotopic to f and makes the diagram below commutative up to homotopy.*

$$\begin{array}{ccc} & & M \times_B M - \Delta \\ & \nearrow_{h_n} & \downarrow i \\ M & \xrightarrow[(1,f^n)]{} & M \times_B M \end{array} \tag{2}$$

Proof. ($\Rightarrow$) Suppose that exists $g : M \to M$, $g \sim_B f$, with $Fix(g^n) = \emptyset$. Is enough to define $h_n = (Id, g^n)$, that is, $s = g$.

($\Leftarrow$) If there is h_n then we have $(Id, f^n) \sim (Id, s^n)$, where $s \sim_B f$. Therefore $Fix(s^n) = \emptyset$. Thus, takes $s = g$. □

3 Torus fiber-preserving maps

Let T be, the torus, defined as the quotient space $\mathbb{R} \times \mathbb{R}/\mathbb{Z} \times \mathbb{Z}$. We denote by (x, y) the elements of $\mathbb{R} \times \mathbb{R}$ and by $[(x, y)]$ the elements in T.

Let $MA = \frac{T\times[0,1]}{([(x,y)],0)\sim(\left[A\binom{x}{y}\right],1)}$ be the quotient space, where A is a homeomorphism of T induced by an operator in $\mathbb{R}^2$ that preserves $\mathbb{Z}\times\mathbb{Z}$. The space MA is a fiber bundle over the circle S^1 where the fiber is the torus. For more details on these bundles see [6].

Given a fiber-preserving map $f : MA \to MA$, i.e. $p \circ f = p$ we want to compute the number $Fix(g^n)$ for each map g fiberwise homotopic to f.

Consider the loops in MA given by; $a(t) =< [(t,0)], 0 >$, $b(t) =< [(0,t)], 0 >$ and $c(t) =< [(0,0)], t >$ for $t \in [0,1]$. We denote by B the matrix of the homomorphism induced on the fundamental group by the restriction of f to the fiber T. From [6] we have the following theorem that provides a relationship between the matrices A and B, where

$$A = \begin{pmatrix} a_1 & a_3 \\ a_2 & a_4 \end{pmatrix}$$

From [6] the induced homomorphism $f_\# : \pi_1(MA) \to \pi_1(MA)$ is given by; $f_\#(a) = a^{b_1}b^{b_2}$, $f_\#(b) = a^{b_3}b^{b_4}$, $f_\#(c) = a^{c_1}b^{c_2}c$. Thus

$$B = \begin{pmatrix} b_1 & b_3 \\ b_2 & b_4 \end{pmatrix}$$

Theorem 3.1. (1) $\pi_1(MA, 0) = \langle a, b, c | [a, b] = 1, cac^{-1} = a^{a_1}b^{a_2}, cbc^{-1} = a^{a_3}b^{a_4}\rangle$

(2) *B commutes with A.*

(3) *If f restricted to the fiber is deformable to a fixed point free map then the determinant of $B - I$ is zero, where I is the identity matrix.*

(4) *If v is an eigenvector of B associated to 1 ($for\, B \neq Id$) then $A(v)$ is also an eigenvector of B associated to 1.*

(5) *Consider $w = A(v)$ if the pair v, w generators $\mathbb{Z}\times\mathbb{Z}$, otherwise let w be another vector so that v, w span $\mathbb{Z}\times\mathbb{Z}$. Define the linear operator $P : \mathbb{R}\times\mathbb{R} \to \mathbb{R}\times\mathbb{R}$ by $P(v) = \binom{1}{0}$ and $P(w) = \binom{0}{1}$. Consider an isomorphism of fiber bundles, also denoted by P, $P : MA \to M(A^1)$ where $A^1 = P\cdot A\cdot P^{-1}$. Then MA is homeomorphic to $M(A^1)$ over S^1. Moreover we have one of the cases of the table below with $B^1 = P \cdot A \cdot P^{-1}$ and $B^1 \neq Id$, except in case I:*

Case I	$A^1 = \begin{pmatrix} a_1 & a_3 \\ a_2 & a_4 \end{pmatrix}$, $B^1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ $a_3 \neq 0$
Case II	$A^1 = \begin{pmatrix} 1 & a_3 \\ 0 & 1 \end{pmatrix}$, $B^1 = \begin{pmatrix} 1 & b_3 \\ 0 & b_4 \end{pmatrix}$ $a_3(b_4 - 1) = 0$
Case III	$A^1 = \begin{pmatrix} 1 & a_3 \\ 0 & -1 \end{pmatrix}$, $B^1 = \begin{pmatrix} 1 & b_3 \\ 0 & b_4 \end{pmatrix}$ $a_3(b_4 - 1) = -2b_3$
Case IV	$A^1 = \begin{pmatrix} -1 & a_3 \\ 0 & -1 \end{pmatrix}$, $B^1 = \begin{pmatrix} 1 & b_3 \\ 0 & b_4 \end{pmatrix}$ $a_3(b_4 - 1) = 0$
Case V	$A^1 = \begin{pmatrix} -1 & a_3 \\ 0 & 1 \end{pmatrix}$, $B^1 = \begin{pmatrix} 1 & b_3 \\ 0 & b_4 \end{pmatrix}$ $a_3(b_4 - 1) = 2b_3$

From [6] we have the following theorem:

Theorem 3.2. *A fiber-preserving map $f : MA \to MA$ can be deformed to a fixed point free map by a homotopy over S^1 if and only if one of the cases below holds:*

(1) MA is as in case I and f is arbitrary

(2) MA is as in one of the cases II or III and $c_1(b_4 - 1) - c_2 b_3 = 0$

(3) MA is as in case IV and $n(b_4(b_3 + 1) - 1 - c_1(b_4 - 1) + b_3 c_2) - (n - 1)(b_4 - 1) \equiv_2 0$ except when:

a_3 is odd and $[(c_1 + b_3 c_2 \frac{(n-1)}{2}, nc_2)] = [(0,0)] \in \frac{\mathbb{Z} \oplus \mathbb{Z}}{\langle (1,2), (0,4) \rangle}$ or

a_3 is even and $[(nc_1 + \frac{n(n-1)}{2} b_3 b_4 c_2, c_2 + (n-1) b_4 c_2)] = [(0,0)]$, with $[(0,0)] \in \frac{\mathbb{Z} \oplus \mathbb{Z}}{\langle (2,0), (0,2) \rangle}$.

(4) MA is as in case V and either

a_3 is even and $(b_4 - 1)(c_1 - \frac{a_3}{2} c_2 - 1) \equiv 0$ mod 2, except when $c_1 - \frac{a_3}{2} c_2 - 1$ and $\frac{b_4 - 1}{L}$ are odd, or

a_3 is odd and $\frac{b_4 - 1}{2}(1 + c_2) \equiv 0$ mod 2 except when $1 + c_2$ and $\frac{b_4 - 1}{L}$ are odd, where $L := mdc(b_4 - 1, c_2)$.

Given $n \in \mathbb{N}$ we denote the induced homomorphism $f^n_\# : \pi_1(MA) \to \pi_1(MA)$ by; $f_\#(a) = a^{b_{1n}} b^{b_{2n}}$, $f_\#(b) = a^{b_{3n}} b^{b_{4n}}$, $f_\#(c) = a^{c_{1n}} b^{c_{2n}} c$, where $b_{j1} = b_j, j = 1, ..., 4$ and $c_{j1} = c_j, j = 1, 2$. Thus the matrix of the homomorphism

induced on the fundamental group by the restriction of f^n to the fiber T is given by;

$$B_n = \begin{pmatrix} b_{1n} & b_{3n} \\ b_{2n} & b_{4n} \end{pmatrix}$$

where $B_1 = B$ is the matrix of $(f_{|T})_\#$ and $B_n = B^n$. From [8] we have

$$N(h^n) = |L(h^n)| = |det([h_\#]^n - I)|$$

for each map $h : T \to T$ on torus, where $[h_\#]$ is the matrix of induced homomorphism and I is the identity.

Since $(B^n - I) = (B - I)(B^{n-1} + ... + B + I)$ then $det(B^n - I) = det(B - I)det(B^{n-1} + ... + B + I)$. Therefore if $f_{|T}$ is deformable to a fixed point free map then $f^n_{|T}$ is deformable to a fixed point free map.

In the Theorems 3.1 and 3.2 putting f^n in the place of f we will get conditions to f^n. The conditions in Theorem 3.1 to f^n is the same of f but the conditions to f^n in the Theorem 3.2 are different of f and are in the Theorem 4.1.

4 Fixed points of f^n

Given a fiber-preserving map $f : MA \to MA$, if $f \sim_{S^1} g$ then $f^n \sim_{S^1} g^n$. Therefore, if $Fix(g^n) = \emptyset$ then the homomorphism $f^n_\# : \pi_1(M) \to \pi(M)$ satisfies the condition of deformability gives in [6].

Proposition 4.1. *Let $f : MA \to MA$ be a fiber-preserving map, where M is a T-bundle over S^1. Suppose which f restrict to the fiber can be deformed to a fixed point free map. This implies $L(f|_T) = 0$. From Theorem 3.1 we can suppose that the induced homomorphism $f_\# : \pi_1(MA) \to \pi_1(MA)$ is given by; $f_\#(a) = a$, $f_\#(b) = a^{b_3}b^{b_4}$, $f_\#(c) = a^{c_1}b^{c_2}c$. Given $n \in \mathbb{N}$ then from relation $(f_\#)^n = f^n_\#$ we obtain;*

$$f^n_\#(a) = a,$$

$$f^n_\#(b) = a^{b_3 \sum_{i=0}^{n-1} b_4^i} b^{b_4^n},$$

$$f^n_\#(c) = a^{nc_1 + b_3c_2 \sum_{i=0}^{n-2}(n-1-i)b_4^i} b^{c_2 \sum_{i=0}^{n-1} b_4^i} c.$$

Proof. In fact, $f^2_\#(b) = f_\#(a^{b_3}b^{b_4}) = a^{b_3}(a^{b_3}b^{b_4})^{b_4} = a^{b_3 + b_3b_4}b^{b_4^2}$ and $f^2_\#(c) = f_\#(a^{c_1}b^{c_2}c) = a^{c_1}(a^{b_3}b^{b_4})^{c_2}(a^{c_1}b^{c_2}c) = a^{2c_1 + b_3c_2}b^{c_2 + c_2b_4}c$. Suppose that $f^n_\#(b) = a^{b_3 \sum_{i=0}^{n-1} b_4^i} b^{b_4^n}$ and $f^n_\#(c) = a^{nc_1 + b_3c_2 \sum_{i=0}^{n-2}(n-1-i)b_4^i} b^{c_2 \sum_{i=0}^{n-1} b_4^i} c$. Then,

$$\begin{aligned} f^{n+1}_\#(b) &= f_\#(a^{b_3 \sum_{i=0}^{n-1} b_4^i} b^{b_4^n}) &&= a^{b_3 \sum_{i=0}^{n-1} b_4^i}(a^{b_3}b^{b_4})^{b_4^n} \\ &= a^{b_3 \sum_{i=0}^{n-1} b_4^i}(a^{b_3b_4^n}b^{b_4^{n+1}}) &&= a^{b_3 \sum_{i=0}^{n} b_4^i} b^{b_4^{n+1}}; \end{aligned}$$

$$\begin{aligned} f_{\#}^{n+1}(c) &= f_{\#}(a^{nc_1+b_3c_2\sum_{i=0}^{n-2}(n-1-i)b_4^i}b^{c_2\sum_{i=0}^{n-1}b_4^i}c) \\ &= a^{nc_1+b_3c_2\sum_{i=0}^{n-2}(n-1-i)b_4^i}(a^{b_3}b^{b_4})^{c_2\sum_{i=0}^{n-1}b_4^i}(a^{c_1}b^{c_2}c) \\ &= a^{(nc_1+b_3c_2\sum_{i=0}^{n-2}(n-1-i)b_4^i)+(b_3c_2\sum_{i=0}^{n-1}b_4^i)+(c_1)}b^{(c_2\sum_{i=1}^{n}b_4^i)+(c_2)}c \\ &= a^{(n+1)c_1+b_3c_2\sum_{i=0}^{n-1}(n-1-i)b_4^i}b^{c_2\sum_{i=0}^{n}b_4^i}c. \end{aligned}$$

□

Theorem 4.1. *Let $f : MA \to MA$ be a fiber-preserving map, where MA is a T-bundle over S^1. Suppose which f restrict to the fiber can be deformed to a fixed point free map and that the induced homomorphism $f_{\#} : \pi_1(MA) \to \pi_1(MA)$ is given by; $f_{\#}(a) = a$, $f_{\#}(b) = a^{b_3}b^{b_4}$, $f_{\#}(c) = a^{c_1}b^{c_2}c$ as in cases of the Theorem 3.2. Then $f^n : MA \to MA$ can be deformed to a fixed point free map over S^1 if and only if the following conditions are satisfies;*

1) MA is as in case I and f is arbitrary.

2) MA is as in cases II, III and $c_1(b_4 - 1) - c_2b_3 = 0$ or n even and $b_4 = -1$.

3) MA is as in case IV and $n(b_4(b_3 + 1) - 1 - c_1(b_4 - 1) + b_3c_2) - (n - 1)(b_4 - 1) \equiv_2 0$ except when:
a_3 is odd and $[(nc_1 + \frac{n(n-1)}{2}b_3c_2, nc_2)] = [(0, 0)] \in \frac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(1,2),(0,4)\rangle}$ or
a_3 is even and $[(nc_1 + \frac{n(n-1)}{2}b_3b_4c_2, c_2 + (n - 1)b_4c_2)] = [(0, 0)] \in \frac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(2,0),(0,2)\rangle}$.

4) MA is as in case V and either
a_3 is even and $n(b_4-1)(c_1-\frac{a_3}{2}c_2-1)+(n-1)(b_4-1) \equiv 0$ mod 2, except when $n(c_1 - \frac{a_3}{2}c_2 - 1) + (n - 1)$ and $\frac{b_4-1}{L}$ are odd, or
a_3 is odd and $\frac{b_4-1}{2}((1 + c_2)(1 + (n - 1)b_4)) \equiv 0$ mod 2 except when $(1 + c_2)(1 + (n - 1)b_4)$ and $\frac{b_4-1}{L}$ are odd, where $L := mdc(b_4 - 1, c_2)$.

Proof. In proof of this theorem we will denote $f_{\#}^n(b) = a^{b'_3}b^{b'_4}$ and $f_{\#}^n(c) = a^{c'_1}b^{c'_2}c$.

(1) From Theorem 3.2 each map $f : MA \to MA$ is fiberwise homotopic to a fixed point free map over S^1 in particular that happens with $f^n : MA \to MA$ for each $n \in \mathbb{N}$.

(2) If $b_4 = 1$ then $b'_3 = nb_3$, $b'_4 = 1$, $c'_1 = nc_1 + b_3c_2\dfrac{n(n-1)}{2}$ and $c'_2 = nc_2$. In this sense, following Theorem 3.2 of [6], in cases II and III, f^n can be deformed, by a fiberwise homotopy, to a fixed point free map if and only if $c'_1(b'_4 - 1) - c'_2b'_3 = 0$. However, $c'_1(b'_4 - 1) - c'_2b'_3 = n^2c_2b_3$, and $n^2c_2b_3 = 0$ if and only if $c_2 = 0$ or $b_3 = 0$.

For $b_4 \neq 1$ we have $b'_3 = b_3\sum_{i=0}^{n-1} b_4^i = b_3\dfrac{b_4^n-1}{b_4-1}$, $b'_4 = b_4^n$, $c'_1 = nc_1 + b_3c_2\sum_{i=0}^{n-2}(n-1-i)b_4^i = nc_1+b_3c_2\dfrac{b_4^n-nb_4+n-1}{(b_4-1)^2}$ and $c'_2 = c_2\sum_{i=0}^{n-1} b_4^i = c_2\dfrac{b_4^n-1}{b_4-1}$. Then, $c'_1(b'_4-1) - c'_2b'_3 = \dfrac{n(b_4^n-1).(c_1(b_4-1)-c_2b_3)}{b_4-1}$. In fact,

$$\begin{aligned} c'_1(b'_4-1) &= \left(nc_1 + c_2b_3\frac{b_4^n-nb_4+n-1}{(b_4-1)^2}\right)(b_4^n-1) \\ &= nc_1(b_4^n-1) + c_2b_3\left(\frac{(b_4^n-1)-n(b_4-1)}{(b_4-1)^2}\right)(b_4^n-1) \\ &= nc_1(b_4^n-1) + c_2b_3\left(\frac{b_4^n-1}{b_4-1}\right)^2 - nc_2b_3\left(\frac{b_4^n-1}{b_4-1}\right); \\ c'_2b'_3 &= \left(c_2\frac{b_4^n-1}{b_4-1}\right)\left(b_3\frac{b_4^n-1}{b_4-1}\right) = c_2b_3\left(\frac{b_4^n-1}{b_4-1}\right)^2. \end{aligned}$$

Therefore,

$$\begin{aligned} c'_1(b'_4-1) - c'_2b'_3 &= nc_1(b_4^n-1) - nc_2b_3\left(\frac{b_4^n-1}{b_4-1}\right) \\ &= n(b_4^n-1)\left(c_1 - \frac{c_2b_3}{b_4-1}\right) \\ &= n\left(\frac{b_4^n-1}{b_4-1}\right)(c_1(b_4-1)-c_2b_3) \\ &= n(c_1(b_4-1)-c_2b_3)\left(\sum_{i=0}^{n-1} b_4^i\right). \end{aligned}$$

Note that $cos\left(\dfrac{2k\pi}{n}\right) + i.sin\left(\dfrac{2k\pi}{n}\right)$, for $k = 0, 1, \ldots, n-1$, are the roots of $b_4^n - 1 = 0$. So, $+1$ and -1 are the only two possible integer solutions $b_4^n - 1 = 0$. Since $b_4^n - 1 \neq 0$ for n odd and $b_4 \neq 1$, we may assume that $\dfrac{n(b_4^n-1).(c_1(b_4-1)-c_2b_3)}{b_4-1} = 0$ if and only if $c_1(b_4-1) - c_2b_3 = 0$. Then, by Theorem 3.2 again, in cases II and III and n odd, f^n can be deformed, by a fiberwise homotopy, to a fixed point free map if and only if f can be deformed, by a fiberwise homotopy, to a fixed point free map. For n even and $b_4 \neq 1$, $b_4^n - 1 = 0$ if and only if $b_4 = -1$.

(3) Following Theorem 3.2 of [6], in cases IV, f^n can be deformed, by a fiberwise homotopy, to a fixed point free map iff $b'_4(b'_3+1) - 1 - c'_1(b'_4-1) +$

$c_2'b_3' \equiv_2 0$ except when a_3 even and $[(c_1', c_2')] = [(0,0)] \in \dfrac{\mathbb{Z}\oplus\mathbb{Z}}{\langle (2,0),(0,2)\rangle}$, or a_3 odd and $[(c_1', c_2')] = [(0,0)] \in \dfrac{\mathbb{Z}\oplus\mathbb{Z}}{\langle (1,2),(0,4)\rangle}$.

As in case (2), we have $-c_1'(b_4'-1)+c_2'b_3' = -n(c_1(b_4-1)-c_2b_3)\left(\sum_{i=0}^{n-1} b_4^i\right)$ and $b_4'(b_3'+1)-1 = b_4^n\left(1+b_3\sum_{i=0}^{n-1} b_4^i\right)-1 = (b_4^n-1)+b_4^n b_3\left(\sum_{i=0}^{n-1} b_4^i\right) = (b_4-1)\left(\sum_{i=0}^{n-1} b_4^i\right)+b_4^n b_3\left(\sum_{i=0}^{n-1} b_4^i\right)$. Then

$$\begin{gathered}
b_4'(b_3'+1)-1-c_1'(b_4'-1)+c_2'b_3' \;=\\
(b_4-1)\left(\sum_{i=0}^{n-1} b_4^i\right)+b_3b_4^n\left(\sum_{i=0}^{n-1} b_4^i\right)-n(c_1(b_4-1)-c_2b_3)\left(\sum_{i=0}^{n-1} b_4^i\right) \;=\\
(b_4^n-1)(1-nc_1)+b_3(b_4^n+nc_2)\left(\sum_{i=0}^{n-1} b_4^i\right) \;\equiv_2\\
(b_4-1)(1-nc_1)+b_3(b_4+nc_2)(1+(n-1)b_4) \;=\\
(b_4-1)(1-nc_1)+b_3(b_4+(n-1)b_4^2+nc_2+n^2b_4c_2-nb_4c_2) \;\equiv_2\\
(b_4-1)(1-nc_1)+b_3(b_4+(n-1)b_4+nc_2+nb_4c_2-nb_4c_2) \;=\\
(b_4-1)(1-nc_1)+b_3(nb_4+nc_2) \;=\\
n(b_4(b_3+1)-1-c_1(b_4-1)+b_3c_2)-(n-1)(b_4-1).
\end{gathered}$$

The exceptions holds for a_3 even and $[(c_1', c_2')] = [(0,0)] \in \dfrac{\mathbb{Z}\oplus\mathbb{Z}}{\langle (2,0),(0,2)\rangle}$, or a_3 odd and $[(c_1', c_2')] = [(0,0)] \in \dfrac{\mathbb{Z}\oplus\mathbb{Z}}{\langle (1,2),(0,4)\rangle}$. In this sense, we have

$$(c_1', c_2') = \left(nc_1 + b_3c_2\sum_{i=0}^{n-1}(n-1-i)b_4^i, c_2\sum_{i=0}^{n-1} b_4^i\right).$$

If a_3 is odd then $b_4 = 1$, $c_2\sum_{i=0}^{n-1} 1^i = nc_2$ and $nc_1 + b_3c_2\sum_{i=0}^{n-1}(n-1-i)1^i = nc_1 + b_3c_2\dfrac{n(n-1)}{2}$. If a_3 is even then $c_2\sum_{i=0}^{n-1} b_4^i \equiv_2 c_2(1+(n-1)b_4)$ and $nc_1 + b_3c_2\sum_{i=0}^{n-1}(n-1-i)b_4^i \equiv_2 nc_1 + \dfrac{n(n-1)}{2}b_3b_4c_2$.

(4) From Theorem 3.2 the map f^n can be deformed, over S^1, to a fixed point free map if and only if the following condition is satisfy;

a_3 is even and $(b_4^{'}-1)(c_1^{'}-\frac{a_3}{2}c_2^{'}-1)\equiv 0\ mod\ 2$, except when $c_1^{'}-\frac{a_3}{2}c_2^{'}-1$ and $\frac{b_4^{'}-1}{L}$ are odd, or a_3 is odd and $\frac{b_4^{'}-1}{2}(1+c_2^{'})\equiv 0\ mod\ 2$ except when $1+c_2^{'}$ and $\frac{b_4^{'}-1}{L}$ are odd, where $L:=mdc(b_4^{'}-1,c_2^{'})$.

Note that if $b_4=1$ then from Theorem 3.1 we must have $b_3=0$ and this situation return in the case I. Therefore let us suppose $b_4\neq 1$.

From previous calculation we have; $b_4^{'}=b_4^n$, $b_3^{'}=b_3\frac{b_4^n-1}{b_4-1}$, $c_2^{'}=c_2\frac{b_4^n-1}{b_4-1}$ and $c_1^{'}=nc_1+b_3c_2\frac{b_4^n-nb_4+n-1}{(b_4-1)^2}$. From Theorem 3.1 we have $a_3(b_4-1)=2b_3$.

Suppose a_3 even. Since $c_1^{'}(b_4^{'}-1)-c_2^{'}b_3^{'}=\frac{n(b_4^n-1)(c_1(b_4-1)-c_2b_3)}{b_4-1}$ Then $(b_4^{'}-1)(c_1^{'}-\frac{a_3}{2}c_2^{'}-1)=n(b_4^n-1)(c_1-\frac{a_3}{2}c_2-1)+(n-1)(b_4^n-1)$. In fact,

$$\begin{aligned} c_1^{'}-\tfrac{a_3}{2}c_2^{'} &= nc_1+b_3c_2\tfrac{b_4^n-nb_4+n-1}{(b_4-1)^2}-\tfrac{a_3}{2}c_2\tfrac{b_4^n-1}{b_4-1} \\ &= nc_1+b_3c_2\tfrac{(b_4^n-1)-n(b_4-1)}{(b_4-1)^2}-b_3c_2\tfrac{b_4^n-1}{(b_4-1)^2} \\ &= nc_1-\tfrac{b_3c_2n}{b_4-1} \\ &= n(c_1-\tfrac{a_3}{2}c_2). \end{aligned}$$

We know that if $L:=mdc(b_4-1,c_2)$ then $kL:=mdc(k(b_4-1),kc_2)$. Thus, $mdc(b_4^{'}-1,c_2^{'})=L^{'}=\frac{b_4^n-1}{(b_4-1)}L=$ where $L=mdc(b_4-1,c_2)$ because $b_4^{'}-1=\frac{b_4^n-1}{(b_4-1)}(b_4-1)$ and $c_2^{'}=c_2\frac{b_4^n-1}{(b_4-1)}$. Furthermore, $\frac{b_4^{'}-1}{L^{'}}=\frac{b_4^{'}-1}{L}\frac{b_4-1}{(b_4^n-1)}=\frac{b_4-1}{L}$. With these calculations we obtain the conditions statements on the theorem.

In the case a_3 odd we must have; $\frac{b_4^n-1}{2}(1+c_2\frac{b_4^n-1}{b_4-1})\equiv 0\ mod\ 2$ except when $1+c_2\frac{b_4^n-1}{b_4-1}$ and $\frac{b_4-1}{L}$ are odd, where $L:=mdc(b_4-1,c_2)$.

Note that $\frac{b_4^n-1}{b_4-1}$ is even if and only if $1+(n-1)b_4$ is even, and b_4^n-1 is even if and only if b_4-1 is even, for all $n\in\mathbb{N}$. With this we obtain the enunciate of the theorem. □

Corollary 4.1. *From Theorem 4.1, if $f:MA\to MA$ is deformed to a fixed point free map over S^1 and $n\in\mathbb{N}$ is odd then the map $f^n:MA\to MA$ can be deformed to a fixed point free map over S^1.*

Proof. If $f:MA\to MA$ is deformed to a fixed point free map over S^1 then the conditions of the Theorem 3.2 are satisfied. Suppose n odd then the conditions of the Theorem 4.1 also are satisfied. Thus $f^n:MA\to MA$ can be deformed to a fixed point free map over S^1. □

In the corollary above if n is even the above statement may not holds, for example in the case V of the Theorem 4.1 if n, b_4, a_3 and $c_1-\frac{a_3}{2}c_2-1$ are even then $f:MA\to MA$ is deformed to a fixed point free map over S^1 but f^n is not.

Proposition 4.2. *Let $f : MA \to MA$ be a fiber-preserving such that the induced homomorphism $f_\# : \pi_1(MA) \to \pi_1(MA)$ is given by; $f_\#(a) = a$, $f_\#(b) = a^{b_3}b^{b_4}$, $f_\#(c) = a^{c_1}b^{c_2}c$. Suppose that for some n odd, $n \in \mathbb{N}$ the fiber-preserving map $f^n : MA \to MA$ is deformed to a fixed point free map over S^1. If k is a divisor of n then the map $f^k : MA \to MA$ can be deformed by a fiberwise homotopy to a fixed point free map over S^1.*

Proof. Is enough to verify that if the conditions of the Theorem 4.1 are satisfied for some n odd then those conditions are also satisfied for any k divisor of n. We will analyze each case of the Theorem 4.1.

Case I. In this case for each $n \in \mathbb{N}$ the fiber-preserving map can be deformed over S^1 to a fixed point free map.

Cases II and III. In these cases if for some n odd the fiber-preserving map $f^n : MA \to MA$ is deformed to a fixed point free map over S^1 then we must have; $c_1(b_4 - 1) - c_2b_3 = 0$. Thus for all $k \leq n$ the f^k can be deformed to a fixed point free map over S^1, in particular when k divides n.

Case IV. Suppose that for some odd positive integer n the fiber-preserving map $f^n : MA \to MA$ is deformed to a fixed point free map over S^1, then $n(b_4(b_3+1) - 1 - c_1(b_4-1) + b_3c_2) - (n-1)(b_4-1) \equiv_2 0$ and
if a_3 is odd then $[(nc_1 + \frac{n(n-1)}{2}b_3c_2, nc_2)] \neq [(0,0)] \in \frac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(1,2),(0,4)\rangle}$ or
if a_3 is even then $[(nc_1 + \frac{n(n-1)}{2}b_3b_4c_2, c_2 + (n-1)b_4c_2)] \neq [(0,0)] \in \frac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(2,0),(0,2)\rangle}$.
Note that if a_3 is odd then $b_4 = 1$ and

$$n(b_4(b_3+1) - 1 - c_1(b_4-1) + b_3c_2) - (n-1)(b_4-1) \equiv_2 b_3 + b_3c_2 = b_3(1+c_2);$$

$$\begin{array}{rcl} [(nc_1 + \frac{n(n-1)}{2}b_3c_2, nc_2)] & = & [(0, nc_2 - 2(nc_1 + \frac{n(n-1)}{2}b_3c_2))] \\ & = & [(0, n(c_2 - 2c_1 - (n-1)b_3c_2)))] \in \frac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(1,2),(0,4)\rangle}; \\ & \Rightarrow & n(c_2 - 2c_1 - (n-1)b_3c_2) \not\equiv_4 0 \\ & \Rightarrow & c_2 - 2c_1 - (n-1)b_3c_2 \not\equiv_4 0. \end{array}$$

If b_3 is even then $c_2 - 2c_1 - (n-1)b_3c_2 \equiv_4 c_2 - 2c_1$. If c_2 is odd then $c_2 - 2c_1 - (n-1)b_3c_2$ is odd and $c_2 - 2c_1 - (k-1)b_3c_2$ is odd for each odd k. If a_3 is even we have

$$\left[\left(nc_1 + \frac{n(n-1)}{2}b_3b_4c_2, c_2 + (n-1)b_4c_2\right)\right] = \left[\left(c_1 + \frac{(n-1)}{2}b_3b_4c_2, c_2\right)\right].$$

Then, $c_2 \equiv_2 1$ or $c_1 + \frac{(n-1)}{2}b_3b_4c_2 \equiv_2 1$. So, if $c_2 \equiv_2 0$ then $c_1 + \frac{(n-1)}{2}b_3b_4c_2 \equiv_2 c_1$. Therefore, $c_1 \equiv_2 1$ or $c_2 \equiv_2 1$.

Let k be an integer such that k divides n then k must be odd and

$$\begin{array}{rcl} k(b_4(b_3+1)-1-c_1(b_4-1)+b_3c_2)-(k-1)(b_4-1) & \equiv_2 & \\ n(b_4(b_3+1)-1-c_1(b_4-1)+b_3c_2)-(n-1)(b_4-1) & \equiv_2 & 0. \end{array}$$

If a_3 is odd then b_3 is even or c_2 is odd, $b_4 = 1$ and

$$\begin{array}{rcl} [(kc_1+\frac{k(k-1)}{2}b_3c_2, kc_2)] & = & [(0, k(c_2-2c_1-(k-1)b_3c_2)))] \\ & = & [(0, c_2-2c_1-(k-1)b_3c_2)))] \\ & = & [(0, c_2-2c_1-(n-1)b_3c_2)))] \\ & \neq & [(0,0)] \in \frac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(1,2),(0,4)\rangle}. \end{array}$$

Then, $f^k : MA \to MA$ can be deformed to a fixed point free map over S^1.

If a_3 is even then

$$\begin{array}{rcl} [(kc_1+\frac{k(k-1)}{2}b_3b_4c_2, c_2+(k-1)b_4c_2)] & = & [(c_1+\frac{(k-1)}{2}b_3b_4c_2, c_2)] \\ & \neq & [(0,0)] \in \frac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(2,0),(0,2)\rangle}. \end{array}$$

Then, $f^k : MA \to MA$ can be deformed to a fixed point free map over S^1.

Case V. Suppose that for some n odd, $n \in \mathbb{N}$ the fiber-preserving map $f^n : MA \to MA$ is deformed to a fixed point free map over S^1. If k divides n then there is $l \in \mathbb{N}$ such that $kl = n$, in particular l must be odd. The conditions to deform f^n in this case with a_3 even are;

a_3 is even and $n(b_4-1)(c_1-\frac{a_3}{2}c_2-1)+(n-1)(b_4-1) \equiv 0 \ mod\ 2$, except when $n(c_1-\frac{a_3}{2}c_2-1)+(n-1)$ and $\frac{b_4-1}{L}$ are odd, where $L := mdc(b_4-1, c_2)$.

We have; $l[k(b_4-1)(c_1-\frac{a_3}{2}c_2-1)+(k-1)(b_4-1)] = lk(b_4-1)(c_1-\frac{a_3}{2}c_2-1)+(lk-l)(b_4-1) = n(b_4-1)(c_1-\frac{a_3}{2}c_2-1)+(n-1)(b_4-1)+(1-l)(b_4-1)$.

From hypothesis $n(b_4-1)(c_1-\frac{a_3}{2}c_2-1)+(n-1)(b_4-1)$ is even. Since l is odd then $(1-l)(b_4-1)$ is even. Therefore $l[k(b_4-1)(c_1-\frac{a_3}{2}c_2-1)+(k-1)(b_4-1)]$ is even. Since l is odd then we must have $k(b_4-1)(c_1-\frac{a_3}{2}c_2-1)+(k-1)(b_4-1) \equiv 0 \ mod\ 2$.

Note that $l[k(c_1-\frac{a_3}{2}c_2-1)+(k-1)] = [n(c_1-\frac{a_3}{2}c_2-1)+(n-1)]+(1-l)$. Thus $l[k(c_1-\frac{a_3}{2}c_2-1)+(k-1)]$ is odd because $[n(c_1-\frac{a_3}{2}c_2-1)+(n-1)]$ is odd by hypothesis and $(1-l)$ is even. Since l is odd then we must have $k(c_1-\frac{a_3}{2}c_2-1)+(k-1)$ odd. Therefore the conditions to deform f^k to a fixed point free map over S^1, in the Theorem 4.1, are satisfied. The case a_3 odd is analogous. □

Proposition 4.3. *Let $f : MA \to MA$ be a fiber-preserving. If m, n are odd, $m, n \in \mathbb{N}$, then f^m is deformable to a fixed point free map over S^1 if and only if f^n is deformable to a fixed point free map over S^1.*

Proof. If m, n are odd and f^m is deformable to a fixed point free map over S^1 then by Proposition 4.2 f is deformable to a fixed point free map over S^1. From Corollary 4.1 f^n is deformable to a fixed point free map over S^1. □

We have a analogous result to n even;

Proposition 4.4. *Let $f : MA \to MA$ be a fiber-preserving map, where MA is a T-bundle over S^1. Suppose which f restrict to the fiber can be deformed to a fixed point free map and that the induced homomorphism $f_\# : \pi_1(MA) \to \pi_1(MA)$ is given by; $f_\#(a) = a$, $f_\#(b) = a^{b_3}b^{b_4}$, $f_\#(c) = a^{c_1}b^{c_2}c$ as in cases of the Theorem 3.2. Given an even positive integer n such that f^n is deformable to a fixed point free map over S^1 then f^k is deformable to a fixed point free map over S^1, for all even positive integer k, except when MA is as in case IV and*
a_3 is odd and $k \equiv_4 0$ or
a_3 is even, $k \equiv_4 0$ and $b_3b_4c_2 \equiv_2 1$.

Proof. Is enough to verify that if the conditions of the Theorem 4.1 are satisfied for some n even then those conditions are also satisfied by every even k. We will analyze each case of the Theorem 4.1.

Case I. In this case for each $n \in \mathbb{N}$ the fiber-preserving map can be deformed over S^1 to a fixed point free map.

Cases II and III. In these cases if for some n odd the fiber-preserving map $f^n : MA \to MA$ is deformed to a fixed point free map over S^1 then we must have; $c_1(b_4 - 1) - c_2b_3 = 0$ or $b_4 = -1$. Thus, for all even k, f^k can be deformed to a fixed point free map over S^1.

Case IV. If n is an even positive integer and $f^n : MA \to MA$ is deformed to a fixed point free map over S^1, then $n(b_4(b_3 + 1) - 1 - c_1(b_4 - 1) + b_3c_2) - (n-1)(b_4 - 1) \equiv_2 0$ and
if a_3 is odd then $[(nc_1 + \frac{n(n-1)}{2}b_3c_2, nc_2)] \neq [(0,0)] \in \frac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(1,2),(0,4)\rangle}$ or
if a_3 is even then $[(nc_1 + \frac{n(n-1)}{2}b_3b_4c_2, c_2 + (n-1)b_4c_2)] \neq [(0,0)] \in \frac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(2,0),(0,2)\rangle}$.
Note that if a_3 is odd then $b_4 = 1$ and

$$\begin{aligned}
[(nc_1 + \tfrac{n(n-1)}{2}b_3c_2, nc_2)] &= [(0, nc_2 - 2(nc_1 + \tfrac{n(n-1)}{2}b_3c_2))] \\
&= [(0, n(c_2 - 2c_1 - (n-1)b_3c_2)))] \in \tfrac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(1,2),(0,4)\rangle}; \\
&\Rightarrow\ n(c_2 - 2c_1 - (n-1)b_3c_2) \not\equiv_4 0 \\
&\Rightarrow\ c_2 - 2c_1 - (n-1)b_3c_2 \equiv_2 1 \text{ and } n \equiv_4 2.
\end{aligned}$$

If a_3 is even we have

$$\left[\left(nc_1+\frac{n(n-1)}{2}b_3b_4c_2, c_2+(n-1)b_4c_2\right)\right]=\left[\left(\frac{n}{2}b_3b_4c_2, c_2(1+b_4)\right)\right].$$

Let k be an even positive integer then k must be odd and

$$k(b_4(b_3+1)-1-c_1(b_4-1)+b_3c_2)-(k-1)(b_4-1) \equiv_2 0$$

Then, $f^k : MA \to MA$ can be deformed to a fixed point free map over S^1 except when:
a_3 is odd and

$$\begin{aligned}[(kc_1+\tfrac{k(k-1)}{2}b_3c_2, kc_2)] &= [(0, k(c_2-2c_1-(k-1)b_3c_2)))] \\ &= [(0,k)))] \\ &= [(0,0)] \in \tfrac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(1,2),(0,4)\rangle}.\end{aligned}$$

or a_3 is even and

$$\begin{aligned}[(kc_1+\tfrac{k(k-1)}{2}b_3b_4c_2, c_2+(k-1)b_4c_2)] &= \left[\left(\tfrac{k}{2}b_3b_4c_2, c_2(1+b_4)\right)\right] \\ &= [(0,0)] \in \tfrac{\mathbb{Z}\oplus\mathbb{Z}}{\langle(2,0),(0,2)\rangle}.\end{aligned}$$

Note that, if f^n can be deformed to a fixed point free map over S^1 but f^k does not then we have $b_3b_4c_2 \equiv_2 1$ and $k \equiv_4 0$.

Case V. If n is an even positive integer and $f^n : MA \to MA$ is deformed to a fixed point free map over S^1, then
if a_3 is odd then $\frac{b_4-1}{2}((1+c_2)(1+(n-1)b_4)) \equiv_2 0$ and at least one of $(1+c_2)(1+(n-1)b_4)$ and $\frac{b_4-1}{L}$ is even, where $L := mdc(b_4-1, c_2)$, or
if a_3 is even then $n(b_4-1)(c_1-\frac{a_3}{2}c_2-1)+(n-1)(b_4-1) \equiv_2 0$ and at least one of $n(c_1-\frac{a_3}{2}c_2-1)+(n-1)$ and $\frac{b_4-1}{L}$ is even, where $L := mdc(b_4-1, c_2)$.

Let a_3 odd and k an even positive integer then

$$\begin{aligned}(1+(k-1)b_4)) &\equiv_2 (1+(n-1)b_4)) \\ \Rightarrow \quad \tfrac{b_4-1}{2}((1+c_2)(1+(k-1)b_4)) &\equiv_2 \tfrac{b_4-1}{2}((1+c_2)(1+(n-1)b_4)) \\ &\equiv_2 0; \\ (1+c_2)(1+(k-1)b_4) &\equiv_2 (1+c_2)(1+(n-1)b_4).\end{aligned}$$

Then, $f^k : MA \to MA$ can be deformed to a fixed point free map over S^1 for a_3 odd. Let a_3 even and k an even positive integer then

$$\begin{aligned}n(b_4-1)(c_1-\tfrac{a_3}{2}c_2-1)+(n-1)(b_4-1) &\equiv_2 b_4-1; \\ n(c_1-\tfrac{a_3}{2}c_2-1)+(n-1) &\equiv_2 1; \\ \Rightarrow \quad k(b_4-1)(c_1-\tfrac{a_3}{2}c_2-1)+(k-1)(b_4-1) &\equiv_2 0.\end{aligned}$$

Then, $f^k : MA \to MA$ can be deformed to a fixed point free map over S^1 for a_3 even.

□

Given $n \in \mathbb{N}$ and $f : MA \to MA$ a fiber-preserving then from Propositions 4.3 and 4.4 the conditions to deform f and f^2 to a fixed point free map over S^1 are enough to deform f^k to a fixed point free map over S^1 for all k divisor of n.

Theorem 4.2. *Let $f : T \times I \to T \times I$ be the map defined by;*

$$f(x, y, t) = (x + b_3 y + c_1 t + \varepsilon, b_4 y + c_2 t + \delta, t).$$

Denoting $f^n(x, y, t) = (x_n, y_n, t)$ then x_n and y_n are given by

$$\begin{aligned} x_n &= x + b_3 y \sum_{i=0}^{n-1} b_4^i + (nc_1 + b_3 c_2 \sum_{i=0}^{n-1} i b_4^{n-1-i}) t + b_3 \delta \sum_{i=0}^{n-1} i b_4^{n-1-i} + n\varepsilon \\ y_n &= b_4^n y + c_2 t \sum_{i=0}^{n-1} b_4^i + \delta \sum_{i=0}^{n-1} b_4^i, \end{aligned}$$

and for each $n \in \mathbb{N}$ and ϵ, δ appropriates the map f^n induces a fiber-preserving map in the fiber bundle MA, as in Theorem 3.1, which we will represent by $f^n(< x, y, t >) =< x_n, y_n, t >$, such that the induces homomorphism $(f^n)_\#$ is as in the Proposition 4.1. Note that the induce homomorphism $f_\#$ is given by; $f_\#(a) = a$, $f_\#(b) = a^{b_3} b^{b_4}$, $f_\#(c) = a^{c_1} b^{c_2} c$.

Proof. Denote $f^n(x, y, t) = (x_n, y_n, t)$ for each $n \in \mathbb{N}$. We have

$$\begin{aligned} x_2 &= x_1 + b_3 y_1 + c_1 t + \varepsilon \\ &= (x + b_3 y + c_1 t + \varepsilon) + b_3(b_4 y + c_2 t + \delta) + c_1 t + \varepsilon \\ &= x + b_3 y(b_4 + 1) + (2c_1 + b_3 c_2) t + b_3 \delta + 2\varepsilon; \\ y_2 &= b_4 y_1 + c_2 t + \delta \\ &= b_4(b_4 y + c_2 t + \delta) + c_2 t + \delta \\ &= b_4^2 y + c_2(b_4 + 1) t + (b_4 + 1)\delta. \end{aligned}$$

Suppose that $f^n(x, y, t) = (x_n, y_n, t)$ as in hypothesis, then

$$f^{n+1}(x, y, t) = (x_n + b_3 y_n + c_1 t + \varepsilon, b_4 y_n + c_2 t + \delta, t) = (x_{n+1}, y_{n+1}, t),$$

where

$$\begin{aligned}
x_{n+1} &= x_n + b_3y_n + c_1t + \varepsilon \\
&= (x + b_3y\sum_{i=0}^{n-1} b_4^i + (nc_1 + b_3c_2\sum_{i=0}^{n-1} ib_4^{n-1-i})t + b_3\delta\sum_{i=0}^{n-1} ib_4^{n-1-i} + n\varepsilon)+ \\
&\quad +b_3(b_4^n y + c_2t\sum_{i=0}^{n-1} b_4^i + \delta\sum_{i=0}^{n-1} b_4^i) + c_1t + \varepsilon \\
&= x + (b_3y\sum_{i=0}^{n-1} b_4^i + b_3yb_4^n) + ((nc_1 + b_3c_2\sum_{i=0}^{n-1} ib_4^{n-1-i})t + c_1t + \\
&\quad b_3c_2t\sum_{i=0}^{n-1} b_4^i)+(b_3\delta\sum_{i=0}^{n-1} ib_4^{n-1-i} + b_3\delta\sum_{i=0}^{n-1} b_4^i) + (n\varepsilon + \varepsilon) \\
&= x + b_3y\sum_{i=0}^{n} b_4^i + ((n+1)c_1 + b_3c_2\sum_{i=0}^{n} ib_4^{n-i})t + b_3\delta\sum_{i=0}^{n} ib_4^{n-i} + (n+1)\varepsilon;
\end{aligned}$$

$$\begin{aligned}
y_{n+1} &= b_4y_n + c_2t + \delta \\
&= b_4(b_4^n y + c_2t\sum_{i=0}^{n-1} b_4^i + \delta\sum_{i=0}^{n-1} b_4^i) + c_2t + \delta \\
&= b_4^{n+1}y + (c_2t\sum_{i=1}^{n} b_4^i + c_2t) + (\delta\sum_{i=1}^{n} b_4^i + \delta) \\
&= b_4^{n+1}y + c_2t\sum_{i=0}^{n} b_4^i + \delta\sum_{i=0}^{n} b_4^i,
\end{aligned}$$

as we wish. Now, to verify that $f^n(< x, y, 0 >) = f^n(< A\binom{x}{y}, 1 >)$ for each $n \in \mathbb{N}$, in $T \times I$, where T is the torus, firstly we will verify this condition for $n = 1$. We have

$$f < x, y, 0 >=< x + b_3y + \varepsilon, b_4y + \delta, 0 > \quad \textit{and}$$

$f < A\binom{x}{y}, 1 >=< (a_1+a_2b_3)x+(a_3+b_3a_4)y+c_1+\varepsilon, b_4a_2x+b_4a_4y+c_2+\delta, 1 >$

But in MA we have $< x, y, 0 >=< A\binom{x}{y}, 1 >$, that is, $< x, y, 0 >=< a_1x+a_3y, a_2x+a_4y, 1 >$. Now we will analyze each case of the Theorem 3.1.

Case I. In this case we need consider $b_3 = 0$ and $b_4 = 1$. Thus, in MA $f < x, y, 0 >=< x+\epsilon, y+\delta, 0 >= < a_1x+a_3y+a_1\epsilon+a_3\delta, a_2x+a_4y+a_2\epsilon+a_4\delta, 1 >$. Note that, $f < A\binom{x}{y}, 1 >=< a_1x + a_3y + c_1 + \epsilon, a_2x + a_4y + c_2 + \delta, 1 >$. Therefore $f < x, y, 0 >= f < A\binom{x}{y}, 1 >$ if $a_1\epsilon + a_3\delta = \epsilon + k$ and $a_2\epsilon + a_4\delta = \delta + l$ where $k, l \in \mathbb{Z}$.

Case II. In this case we have $a_1 = a_4 = 1$, $a_2 = 0$ and $a_3(b_4 - 1) = 0$. Therefore, $f < A\binom{x}{y}, 1 >=< x + (a_3 + b_3)y + c_1 + \epsilon, b_4y + c_2 + \delta, 1 >$. Thus, $f < x, y, 0 >=< x + b_3y + \epsilon, b_4y + \delta, 0 >=< x + (a_3 + b_3)y + \epsilon + a_3\delta, b_4y + \delta, 1 > =< x + (a_3 + b_3)y + \epsilon + a_3\delta, b_4y + \delta, 1 >$. Therefore $f < x, y, 0 >= f < A\binom{x}{y}, 1 >$ if $a_3\delta \in \mathbb{Z}$.

Case III. In this case we have $a_1 = 1$, $a_4 = -1$, $a_2 = 0$ and $a_3(b_4 - 1) = -2b_3$. Therefore $f < A\binom{x}{y}, 1 >=< x + (a_3 - b_3)y + c_1 + \epsilon, -b_4y + c_2 + \delta, 1 >$. We have $f < x, y, 0 >=< x + b_3y + \epsilon, b_4y + \delta, 0 >=< x + (a_3b_4 + b_3)y + \epsilon + a_3\delta, -b_4y - \delta, 1 >$. Thus $f < x, y, 0 >=< x + (a_3 - b_3)y + \epsilon + a_3\delta, -b_4y - \delta, 1 >$. Then $f < x, y, 0 >= f < A\binom{x}{y}, 1 >$ if $a_3\delta \in \mathbb{Z}$ and $\delta = \frac{k}{2}, k \in \mathbb{Z}$.

Case IV. In this case we have $a_1 = -1$, $a_4 = -1$, $a_2 = 0$ and $a_3(b_4 - 1) = 0$. Thus $f < A\binom{x}{y}, 1 >=< -x + (a_3 - b_3)y + c_1 + \epsilon, -b_4y + c_2 + \delta, 1 >$. We have $f(x, y, 0) = (x + b_3y + \epsilon, b_4y + \delta, 0) = (-x + (a_3b_4 - b_3)y - \epsilon + a_3\delta, -b_4y - \delta, 1)$ Thus $f < x, y, 0 >=< -x + (a_3 - b_3)y - \epsilon + a_3\delta, -b_4y - \delta, 1 >$. Then $f < x, y, 0 >= f < A\binom{x}{y}, 1 >$ if $\epsilon = \frac{a_3m + 2k}{4}$ and $\delta = \frac{m}{2}$ where $m, k \in \mathbb{Z}$.

Case V. In this case we have $a_1 = -1$, $a_4 = 1$, $a_2 = 0$ and $a_3(b_4 - 1) = 2b_3$. Therefore $f < A\binom{x}{y}, 1 >=< -x + (a_3 + b_3)y + c_1 + \epsilon, b_4y + c_2 + \delta, 1 >$. We have $f(x, y, 0) = (x + b_3y + \epsilon, b_4y + \delta, 0) = (-x + (a_3b_4 - b_3)y - \epsilon + a_3\delta, b_4y + \delta, 1)$ Thus $f < x, y, 0 >=< x + (a_3 + b_3)y - \epsilon + a_3\delta, b_4y + \delta, 1 >$. Then $f < x, y, 0 >= f < A\binom{x}{y}, 1 >$ if $\epsilon = \frac{a_3\delta + k}{2}$ where $k \in \mathbb{Z}$.

Now we will verify the condition for all $n \in \mathbb{N}$.

Case I. In MA we have

$$\begin{aligned} f^n < x, y, 0 > &= < x + n\varepsilon, y + n\delta, 0 > \\ &= < a_1(x + n\varepsilon) + a_3(y + n\delta), a_2(x + n\varepsilon) + a_4(y + n\delta), 1 > \\ &= < a_1x + a_3y + na_1\varepsilon + na_3\delta, a_2x + a_4y + na_2\varepsilon + na_4\delta, 1 > . \end{aligned}$$

But, $f^n \left\langle A\binom{x}{y}, 1 \right\rangle =< a_1x + a_3y + nc_1 + n\varepsilon, a_2x + a_4y + nc_2 + n\delta, 1 >$. Then, $f^n < x, y, 0 >= f^n \left\langle A\binom{x}{y}, 1 \right\rangle$ if $na_1\varepsilon + na_3\delta = n\varepsilon + k$, $k \in \mathbb{Z}$, and $na_2\varepsilon + na_4\delta = n\delta + l$, $l \in \mathbb{Z}$.

Case II.

$$
\begin{aligned}
&f^n < x, y, 0 >= \\
&=< x + b_3 y \sum_{i=0}^{n-1} b_4^i + b_3 \delta \sum_{i=0}^{n-1} i b_4^{n-1-i} + n\varepsilon, b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i, 0 > \\
&=< \left(x + b_3 y \sum_{i=0}^{n-1} b_4^i + b_3 \delta \sum_{i=0}^{n-1} i b_4^{n-1-i} + n\varepsilon \right) + a_3 \left(b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i \right), \\
&b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i, 1 > \\
&=< x + \left(a_3 b_4^n + b_3 \sum_{i=0}^{n-1} b_4^i \right) y + \left(a_3 \sum_{i=0}^{n-1} b_4^i + b_3 \sum_{i=0}^{n-1} i b_4^{n-1-i} \right) \delta + n\varepsilon, \\
&b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i, 1 >
\end{aligned}
$$

But, $f^n \left(< A \binom{x}{y}, 1 > \right) =$

$$
\begin{aligned}
&=< x + \left(a_3 + b_3 \sum_{i=0}^{n-1} b_4^i \right) y + b_3 \delta \sum_{i=0}^{n-1} i b_4^{n-1-i} + n\varepsilon + b_3 c_2 \sum_{i=0}^{n-1} i b_4^{n-1-i} + n c_1, \\
&b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i + c_2 \sum_{i=0}^{n-1} b_4^i, 1 > .
\end{aligned}
$$

Thus $f^n \left(< A \binom{x}{y}, 1 > \right) = f^n < x, y, 0 >$ if $\delta a_3 \sum_{i=0}^{n-1} b_4^i \in \mathbb{Z}$.

Case III. $f^n < x, y, 0 >=$

$$
\begin{aligned}
&=< x + b_3 y \sum_{i=0}^{n-1} b_4^i + b_3 \delta \sum_{i=0}^{n-1} i b_4^{n-1-i} + n\varepsilon, b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i, 0 > \\
&=< \left(x + b_3 y \sum_{i=0}^{n-1} b_4^i + b_3 \delta \sum_{i=0}^{n-1} i b_4^{n-1-i} + n\varepsilon \right) + a_3 \left(b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i \right), \\
&-b_4^n y - \delta \sum_{i=0}^{n-1} b_4^i, 1 > \\
&=< x + \left(a_3 b_4^n + b_3 \sum_{i=0}^{n-1} b_4^i \right) y + \left(a_3 \sum_{i=0}^{n-1} b_4^i + b_3 \sum_{i=0}^{n-1} i b_4^{n-1-i} \right) \delta + n\varepsilon, \\
&-b_4^n y - \delta \sum_{i=0}^{n-1} b_4^i, 1 >
\end{aligned}
$$

Note that, $f^n\left(< A\binom{x}{y}, 1 >\right) =$

$$=< x + \left(a_3 - b_3\sum_{i=0}^{n-1} b_4^i\right) y - b_3\delta\sum_{i=0}^{n-1} ib_4^{n-1-i} + n\varepsilon - b_3c_2\sum_{i=0}^{n-1} ib_4^{n-1-i} + nc_1,$$
$$-b_4^n y + \delta\sum_{i=0}^{n-1} b_4^i + c_2\sum_{i=0}^{n-1} b_4^i, 1 > .$$

Therefore, $f^n < x, y, 0 >= f^n\left(< A\binom{x}{y}, 1 >\right)$ if $2\delta\sum_{i=0}^{n-1} b_4^i \in \mathbb{Z}$ and $\left(a_3\sum_{i=0}^{n-1} b_4^i + 2b_3\sum_{i=0}^{n-1} ib_4^{n-1-i}\right)\delta \in \mathbb{Z}$.

Case IV. $f^n < x, y, 0 >=$

$$=< x + b_3y\sum_{i=0}^{n-1} b_4^i + b_3\delta\sum_{i=0}^{n-1} ib_4^{n-1-i} + n\varepsilon, b_4^n y + \delta\sum_{i=0}^{n-1} b_4^i, 0 >$$
$$=< -\left(x + b_3y\sum_{i=0}^{n-1} b_4^i + b_3\delta\sum_{i=0}^{n-1} ib_4^{n-1-i} + n\varepsilon\right) + a_3\left(b_4^n y + \delta\sum_{i=0}^{n-1} b_4^i\right),$$
$$-b_4^n y - \delta\sum_{i=0}^{n-1} b_4^i, 1 >$$

$$=< -x + \left(a_3b_4^n - b_3\sum_{i=0}^{n-1} b_4^i\right) y + \left(a_3\sum_{i=0}^{n-1} b_4^i - b_3\sum_{i=0}^{n-1} ib_4^{n-1-i}\right)\delta - n\varepsilon,$$
$$-b_4^n y - \delta\sum_{i=0}^{n-1} b_4^i, 1 >$$

We have, $f^n\left(< A\binom{x}{y}, 1 >\right) =$

$$=< -x + \left(a_3 - b_3\sum_{i=0}^{n-1} b_4^i\right) y - b_3\delta\sum_{i=0}^{n-1} ib_4^{n-1-i} + n\varepsilon - b_3c_2\sum_{i=0}^{n-1} ib_4^{n-1-i} + nc_1,$$
$$-b_4^n y + \delta\sum_{i=0}^{n-1} b_4^i + c_2\sum_{i=0}^{n-1} b_4^i, 1 >$$

Thus, $f^n < x, y, 0 >= f^n\left(< A\binom{x}{y}, 1 >\right)$ if $2\delta\sum_{i=0}^{n-1} b_4^i \in \mathbb{Z}$ and $2n\varepsilon = a_3\delta\sum_{i=0}^{n-1} b_4^i +$ k, $k \in \mathbb{Z}$.

Case V. $f^n < x, y, 0 >=$

$$=< x + b_3 y \sum_{i=0}^{n-1} b_4^i + b_3 \delta \sum_{i=0}^{n-1} i b_4^{n-1-i} + n\varepsilon, b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i, 0 >$$

$$=< -\left(x + b_3 y \sum_{i=0}^{n-1} b_4^i + b_3 \delta \sum_{i=0}^{n-1} i b_4^{n-1-i} + n\varepsilon \right) + a_3 \left(b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i \right),$$

$$b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i, 1 >$$

$$=< -x + \left(a_3 b_4^n - b_3 \sum_{i=0}^{n-1} b_4^i \right) y + \left(a_3 \sum_{i=0}^{n-1} b_4^i - b_3 \sum_{i=0}^{n-1} i b_4^{n-1-i} \right) \delta - n\varepsilon,$$

$$b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i, 1 >$$

We have, $f^n \left(< A \binom{x}{y}, 1 > \right) =$

$$=< -x + \left(a_3 + b_3 \sum_{i=0}^{n-1} b_4^i \right) y - b_3 \delta \sum_{i=0}^{n-1} i b_4^{n-1-i} + n\varepsilon - b_3 c_2 \sum_{i=0}^{n-1} i b_4^{n-1-i} + n c_1,$$

$$+ b_4^n y + \delta \sum_{i=0}^{n-1} b_4^i + c_2 \sum_{i=0}^{n-1} b_4^i, 1 > .$$

Therefore, $f^n < x, y, 0 >= f^n \left(< A \binom{x}{y}, 1 > \right)$ if $2n\varepsilon = a_3 \delta \sum_{i=0}^{n-1} b_4^i + k$, $k \in \mathbb{Z}$.

Thus for each $n \in \mathbb{N}$ and ϵ, δ satisfying the conditions above the map $f^n : T \times I \to T \times I$ induces a fiber-preserving map on MA which will be represent by the same symbol. □

The next result we will help us to study the fixed points of f^n given in Theorem 4.2.

Proposition 4.5. *Let $n,\ b_3,\ b_4,\ c_1,\ c_2 \in \mathbb{Z}$, $n \geq 1$. If $c_1(b_4 - 1) - c_2 b_3 \neq 0$ then for all $\varepsilon,\ \delta \in \mathbb{R}$ there are $a,\ b \in \mathbb{Z}$ such that the system bellow has solution $(x, y, t) \in \mathbb{R}^2 \times I$:*

$$\begin{cases} x + a &= x + b_3 y \sum_{i=0}^{n-1} b_4^i + (n c_1 + b_3 c_2 \sum_{i=0}^{n-1} i b_4^{n-1-i}) t + b_3 \delta \sum_{i=0}^{n-1} i b_4^{n-1-i} + n\varepsilon; \\ y + b &= b_4^n y + c_2 t \sum_{i=0}^{n-1} b_4^i + \delta \sum_{i=0}^{n-1} b_4^i. \end{cases}$$

Proof: Suppose $b_4 \neq 1$ and $b_4 \neq -1$ with n even ($b_4 = -1$ with n odd is allowed) and $c_1(b_4-1) - b_3c_2 \neq 0$ then given $\varepsilon,\ \delta \in \mathbb{R}$ we have the solutions $x \in \mathbb{R}$ and:

$$\begin{aligned} t &= \frac{nb_3\delta - n(b_4-1)\varepsilon - (b_4-1)a - b_3b}{n(c_1(b_4-1) - b_3c_2)} \in I; \\ y &= \frac{nc_2\varepsilon - nc_1\delta - ac_2}{n(c_1(b_4-1) - b_3c_2)} + b\left(\frac{1}{b_4^n - 1} + \frac{b_3c_2}{n(b_4-1)(c_1(b_4-1) - b_3c_2)}\right) \in \mathbb{R}. \end{aligned}$$

Thus, we need to find $a,\ b \in \mathbb{Z}$ such that $0 \leq t \leq 1$. Let $k_0 = n(c_1(b_4-1) - b_3c_2) \in \mathbb{Z}$, $k_0 \neq 0$, and $k_1 = nb_3\delta - n(b_4-1)\varepsilon \in \mathbb{R}$, $t = \dfrac{k_1 - (b_4-1)a - b_3b}{k_0}$. If $0 \leq k_1 \leq k_0$ or $k_0 \leq k_1 \leq 0$ let $a = b = 0$, then $t = \dfrac{k_1}{k_0}$. If $0 < k_0 \leq k_1$ or $k_1 \leq 0 < k_0$ then there are $d,\ q \in \mathbb{Z}$ such that $k_1 = dk_0 + q$ with $0 \leq q < k_0$. Let $a = nc_1d$ and $b = nc_2d$, then

$$t = \frac{dk_0 + q - (b_4-1)nc_1d - b_3nc_2d}{k_0} = d + \frac{q}{k_0} - \frac{dk_0}{k_0} = \frac{q}{k_0}.$$

If $k_1 \leq k_0 < 0$ or $k_0 < 0 \leq k_1$ then there are $d,\ q \in \mathbb{Z}$ such that $k_1 = dk_0 + q$ with $0 \leq q < |k_0|$. Let $k \in \mathbb{Z}$ the least integer greater than $\dfrac{-q}{k_0}$, $a = nc_1(d-k)$ and $b = nc_2(d-k)$, then

$$t = \frac{dk_0 + q - (b_4-1)nc_1(d-k) - b_3nc_2(d-k)}{k_0} = d + \frac{q}{k_0} - (d-k) = \frac{q}{k_0} + k.$$

Then, $0 \leq t \leq 1$.

If $b_4 = 1$ and $c_1(b_4-1) - b_3c_2 \neq 0$ then $b_3c_2 \neq 0$. Thus, given $\varepsilon,\ \delta \in \mathbb{R}$ we have the solutions $x \in \mathbb{R}$ and:

$$\begin{aligned} t &= \frac{b}{nc_2} - \frac{\delta}{c_2} \in I; \\ y &= \frac{-nc_2\varepsilon + nc_1\delta + ac_2}{nb_3c_2} - b\left(\frac{c_1}{nb_3c_2} + \frac{n-1}{2n}\right) \in \mathbb{R}. \end{aligned}$$

We need to find $b \in \mathbb{Z}$ such that $0 \leq t \leq 1$. If $c_2 > 0$ take $n\delta \leq b \leq n(c_2 + \delta)$ and if $c_2 < 0$ take $n\delta \geq b \geq n(c_2 + \delta)$.

If $b_4 = -1$, n even and $c_1(b_4-1) - b_3c_2 \neq 0$ then $2c_1 + b_3c_2 \neq 0$. Thus, given $\varepsilon,\ \delta \in \mathbb{R}$ we have the solutions $x,\ y \in \mathbb{R}$ and:

$$t = \frac{2a - nb_3\delta - 2n\varepsilon}{n(2c_1 + b_3c_2)} \in I;$$

We need to find $a \in \mathbb{Z}$ such that $0 \leq t \leq 1$. Let $n = 2k$, $k_0 = k(2c_1 + b_3c_2) \in \mathbb{Z}$, $k_0 \neq 0$, and $k_1 = -kb_3\delta - 2k\varepsilon \in \mathbb{R}$, $t = \dfrac{a + k_1}{k_0}$. If $0 \leq k_1 \leq k_0$ or $k_0 \leq k_1 \leq 0$ let $a = 0$, then $t = \dfrac{k_1}{k_0}$. If $0 < k_0 \leq k_1$ or $k_1 \leq 0 < k_0$ then there are $d,\ q \in \mathbb{Z}$ such that $k_1 = dk_0 + q$ with $0 \leq q < k_0$. Let $a = -k_0d$, then $t = \dfrac{-k_0d + dk_0 + q}{k_0} = \dfrac{q}{k_0}$. If $k_1 \leq k_0 < 0$ or $k_0 < 0 \leq k_1$ then there are $d,\ q \in \mathbb{Z}$ such that $k_1 = dk_0 + q$ with $0 \leq q < |k_0|$. Let $l \in \mathbb{Z}$ the greatest integer lower than q and $a = -l - k_0d$, then $t = \dfrac{-l - k_0d + dk_0 + q}{k_0} = \dfrac{q - l}{k_0}$. Then, $0 \leq t \leq 1$.

□

Theorem 4.3 (Main Theorem)**.** *Let $f : MA \to MA$ be a fiber-preserving map, where MA is a T-bundle over S^1, and $f_{\#}(a) = a$, $f_{\#}(b) = a^{b_3}b^{b_4}$, $f_{\#}(c) = a^{c_1}b^{c_2}c$. Suppose that $f^n : MA \to MA$ can be deformed to a fixed point free map over S^1. Then there exits a map g fiberwise homotopic to f such that g^n is a fixed point free map, in the cases I and II, if and only if the following conditions are satisfies;*

1) MA is as in case I and f is arbitrary.
2) MA is as in cases II, III and $c_1(b_4 - 1) - c_2b_3 = 0$.

Proof. (1) For each map f such that $(f_{|T})_{\#} = Id$ consider the map g fiberwise homotopic to f given by; $g'(< x, y, t >) =< x+c_1t+\epsilon, y+c_2t+\delta, t >$, with $\epsilon, \delta \in \mathbb{Q}-\mathbb{Z}$ satisfying the condition; $a_1\epsilon+a_3\delta = \epsilon+k$ and $a_2\epsilon+a_4\delta = \epsilon+k$ for some $k, l \in \mathbb{Z}$. Note that g' is fiberwise homotopic to the map g defined by;

$$g(< x, y, t >) = \begin{cases} < x + 2c_1t + \epsilon, y + \delta, t > & if \quad 0 \leq t \leq \frac{1}{2} \\ < x + c_1 + \epsilon, y + c_2(2t - 1) + \delta, t > & if \quad \frac{1}{2} \leq t \leq 1 \end{cases}$$

In fact, $H : MA \times I \to MA$ defined by;

$$H(< x, y, t >, s) = \begin{cases} < x + c_1t + \epsilon, y + c_2t + \delta, t > & if \quad 0 \leq t \leq s \\ < x + c_1(2t - s) + \epsilon, y + c_2s + \delta, t > & if \quad s \leq t \leq \frac{s+1}{2} \\ < x + c_1 + \epsilon, y + c_2(2t - 1) + \delta, t > & if \quad \frac{s+1}{2} \leq t \leq 1 \end{cases}$$

is a homotopy between g' and g. Note that,

$$g^n(< x, y, t >) = \begin{cases} < x + n2c_1t + n\epsilon, y + n\delta, t > & if \quad 0 \leq t \leq \frac{1}{2} \\ < x + nc_1 + n\epsilon, y + nc_2(2t - 1) + n\delta, t > & if \quad \frac{1}{2} \leq t \leq 1 \end{cases}$$

In this case we have $Fix(g^n) = \emptyset$.

(2) From Theorem 4.1 we must consider two situations; $c_1(b_4-1)-c_2b_3 = 0$ or n even and $b_4 = -1$. First we will suppose $c_1(b_4-1)-c_2b_3 = 0$. Note that if $c_1(b_4-1)-c_2b_3 \neq 0$ then g can not be deformed to a fixed point free map. Therefore can not exits f fiberwise homotopic to g such that $Fix(f^n) = \emptyset$ for all $n \geq 1$ by Lemma 2.1.

Suppose $b_4 = 1$ and $f^n(x,y,t) = (x_n, y_n, t)$ for each $n \in \mathbb{N}$ gives in the Theorem 4.2. Then:

$$\begin{cases} x_n &= x + nb_3y + \left(c_1 + b_3c_2\dfrac{n-1}{2}\right)nt + \dfrac{n(n-1)}{2}b_3\delta + n\varepsilon, \\ y_n &= y + nc_2t + n\delta. \end{cases}$$

For $c_2b_3 = 0$, $f^n(x,y,t)$ is a fixed point free map for each n choosing $\epsilon \in \mathbb{R}-\mathbb{Q}$ and $\delta = 0$ if $b_3 = 0$ or $\delta \in \mathbb{R}-\mathbb{Q}$ if $c_2 = 0$. In fact, if $b_3 = 0$ and $c_2 \neq 0$ choose $\epsilon \in \mathbb{R}-\mathbb{Q}$ and $\delta = 0$, then

$$\begin{cases} x + k_n &= x + nc_1t + n\varepsilon \\ y + l_n &= y + nc_2t \end{cases}$$

for some $k_n, l_n \in \mathbb{Z}$. From equations we obtain $t = \dfrac{l_n}{nc_2} \in \mathbb{Q}$ and $\epsilon = \dfrac{k_n}{n} - \dfrac{c_1l_n}{nc_2} \in \mathbb{Q}$, but this is a contradiction because ϵ in $\mathbb{R}-\mathbb{Q}$. Therefore f^n has no fixed point for all n. If $c_2 = 0$ we choose $\delta \in \mathbb{R}-\mathbb{Q}$ then $y + n\delta = y + l_n$ which implies $\delta = \frac{l_n}{n} \in \mathbb{Q}$ that is a contradiction because δ in $\mathbb{R}-\mathbb{Q}$. Thus f^n has no fixed point for all n.

Now we suppose $b_4 \neq 1$, $\delta = 0$ and $c_1(b_4-1) = c_2b_3$. Then

$$\begin{aligned} x_n &= x + b_3y\sum_{i=0}^{n-1} b_4^i + (nc_1 + b_3c_2\sum_{i=0}^{n-1} ib_4^{n-1-i})t + n\varepsilon \\ &= x + \left(\frac{b_3(b_4^n-1)}{b_4-1}\right)y + \left(nc_1 + \frac{b_3c_2(b_4^n - 1 + n(1-b_4))}{(b_4-1)^2}\right)t + n\varepsilon \\ &= x + \left(\frac{b_4^n-1}{b_4-1}\right)b_3y + \left(\frac{b_4^n-1}{b_4-1}\right)c_1t + n\varepsilon; \\ y_n &= b_4^n y + c_2t\sum_{i=0}^{n-1} b_4^i \\ &= b_4^n y + \left(\frac{c_2(b_4^n-1)}{b_4-1}\right)t. \end{aligned}$$

We are interested at the solutions of the system above:

$$x_n = x + k_n; \tag{3}$$
$$y_n = y + l_n, \tag{4}$$

where $k_n, l_n \in \mathbb{Z}$ and $n \in \mathbb{N}$. With the equation 4 and $c_2 \neq 0$, we have:

$$t = \frac{l_n(b_4 - 1)}{c_2(b_4^n - 1)} + \frac{1 - b_4}{c_2} y.$$

For this t we obtain:

$$\begin{aligned} x_n &= x + \left(\frac{b_4^n - 1}{b_4 - 1}\right) b_3 y + \left(\frac{b_4^n - 1}{b_4 - 1}\right) c_1 t + n\varepsilon \\ &= x - \left(\frac{(b_4^n - 1)((b_4 - 1)c_1 - c_2 b_3)}{(b_4 - 1)c_2}\right) y + n\varepsilon + \frac{c_1 l_n}{c_2} \\ &= x + n\varepsilon + \frac{c_1 l_n}{c_2} \end{aligned}$$

But, for $\varepsilon \in \mathbb{R} - \mathbb{Q}$ the equation 4 has no solution for all $n \in \mathbb{N}$. In fact, for any $k_n \in \mathbb{Z}$ we have:

$$x_n = x + k_n \Rightarrow x + n\varepsilon + \frac{c_1 l_n}{c_2} = x + k_n \Rightarrow \underbrace{\varepsilon}_{\in \mathbb{R} - \mathbb{Q}} = \underbrace{\frac{k_n}{n} - \frac{c_1 l_n}{n c_2}}_{\in \mathbb{Q}}.$$

In the other side, if $c_2 = 0$ then $c_1 = 0$ because $b_4 \neq 1$ and:

$$\begin{aligned} x_n &= x + \left(\frac{b_4^n - 1}{b_4 - 1}\right) b_3 y + \left(\frac{b_4^n - 1}{b_4 - 1}\right) c_1 t + n\varepsilon \\ &= x + \left(\frac{b_4^n - 1}{b_4 - 1}\right) b_3 y + n\varepsilon; \\ y_n &= b_4^n y + \left(\frac{c_2(b_4^n - 1)}{b_4 - 1}\right) t \\ &= b_4^n y. \end{aligned}$$

Thus, $y = \dfrac{l_n}{b_4^n - 1}$ by equation 4. Then, $x_n = x + n\varepsilon + \dfrac{b_3 l_n}{b_4 - 1}$ and the equation 3 has no solution for $\varepsilon \in \mathbb{R} - \mathbb{Q}$ and for $n \in \mathbb{N}$. In fact, for any $k_n, l_n \in \mathbb{Z}$ we have:

$$x_n = x + k_n \Rightarrow x + n\varepsilon + \frac{b_3 l_n}{b_4 - 1} = x + k_n \Rightarrow \underbrace{\varepsilon}_{\in \mathbb{R} - \mathbb{Q}} = \underbrace{\frac{k_n}{n} - \frac{b_3 l_n}{n(b_4 - 1)}}_{\in \mathbb{Q}}.$$

If $b_4 = -1$ and n even then $f^n(x, y, t) = (x_n, y_n, t)$ such that:

$$\begin{aligned} x_n &= x + b_3 \delta\left(\tfrac{n}{2}\right) + n\varepsilon; \\ y_n &= y. \end{aligned}$$

Thus, choosing $\delta = 0$ and $\varepsilon \in \mathbb{R} - \mathbb{Q}$, the first equation has no solution, that is, $Fix(f) = \emptyset$. □

With a similar way of the Theorem 4.3 we obtain the following result.

Theorem 4.4. *Let $f : MA \to MA$ be a fiber-preserving map, where MA is a T-bundle over S^1, and $f_\#(a) = a$, $f_\#(b) = a^{b_3}b^{b_4}$, $f_\#(c) = a^{c_1}b^{c_2}c$. Suppose that $f^n : MA \to MA$ can be deformed to a fixed point free map over S^1 as in the Theorem 4.1. If the following conditions are satisfied, in the cases IV and V, then there exits a map g fiberwise homotopic to f such that g^n is a fixed point free map.*

MA is as in case IV, V and

$$c_1(b_4 - 1) - c_2b_3 = 0 \text{ and } n \text{ odd or}$$

$$c_1(b_4 - 1) - c_2b_3 = 0,\ b_4 \text{ odd and } n = 4k + 2, k \geq 0.$$

References

[1] B.J. Jiang; *Lectures on the Nielsen Fixed Point Theory,* Contemp. Math., vol. 14, Amer. Math. Soc., Providence, 1983.

[2] J. Jezierski; *Weckens theorem for periodic points in dimension at least 3*, Topology and its Applications 153 (2006) 18251837.

[3] J. Jezierski and W. Marzantowicz; *Homotopy Methods in Topological Fixed and Periodic Point Theory*, vol. 3, Topological Fixed Point Theory and Its Applications, Springer, 2006.

[4] E. Fadell and S. Hussein; *A fixed point theory for fibre-preserving maps* Lectures Notes in Mathematics, vol.886, Springer Verlag, 1981, 49-72.

[5] D. L. Gonçalves; *Fixed points of S^1-fibrations,* Pacific J. Math. 129, 1987, 297-306.

[6] D. L. Gonçalves, D.Penteado and J.P Vieira; *Fixed Points on Torus Fiber Bundles over the Circle*, Fundamenta Mathematicae, vol.183 (1), 2004, 1-38.

[7] B. Halpern, *Periodic points on tori*, Pacific J. Math. 83 (1979), no. 1, 117133.

[8] Edward Keppelmann, *Periodics points on nilmanifolds and solvmanifolds*, Pacific Journal of Mathematics, vol.164 (1) (1994), 105–128.

[9] J. W. Vick; *Homology Theory: An Introduction to Algebraic Topology,* Academic Press.

[10] G. W. Whitehead; *Elements of Homotopy Theory,* Springer-Verlag, 1918.